

MRI Enterography With and Without Guideline (62E)

CPT Code: 74183, 72197

Indications: Crohns disease, diarrhea

Contrast: Yes (see MRG MRI Contrast Administration Policy),

Oral Contrast: NPO 6 hours prior to exam

Option 1 – If patient is <125 lbs

1 bottle of Barium Sulfate (e.g. Redicat) given 20 minutes immediately prior to being placed on the table

Option 2 – If patient is >125 lbs

2 bottle of Barium Sulfate (e.g. Redicat) given 40 minutes immediately prior to being placed on the table
(1 bottle at 40min and 1 bottle at 20 min)

Non Contrast:

Sequence	Plane	Slice Thickness	Gap	FOV	Notes
T2 Haste	Coronal	4mm	1	350-450	Fat Sat
True Fisp	CINE 3D	7mm	-5	350-450	
TrueFisp	Coronal	3.5mm	0	350-450	
T2 Haste	Axial Upper	4mm	1	350-450	Fat Sat
T2 Haste	Axial Lower	4mm	1	350-450	Fat Sat
DWI	Axial Upper	7mm	1	350-450	Fat Sat
DWI	Axial Lower	7mm	1	350-450	Fat Sat

*INJECT GLUCAGON IV

Sequence	Plane	Slice Thickness	Gap	FOV	Notes
T1 VIBE 3D	Coronal	2mm	0	350-450	Fat Sat

Contrast:

Sequence	Plane	Slice Thickness	Gap	FOV	Notes
T1 VIBE 3D	Coronal	2mm	0	350-450	Fat Sat, 20 sec, 90 sec, 2.5 min
T1 VIBE 3D	Axial	2mm	0	350-450	Fat Sat